

THE “AMPLITUDE” PARAMETER OF GAMMA-RAY BURSTS AND ITS IMPLICATIONS FOR GRB CLASSIFICATION

HOU-JUN LÜ¹, BING ZHANG¹, EN-WEI LIANG^{2,1}, BIN-BIN ZHANG³, AND TAKANORI SAKAMOTO⁴

ABSTRACT

Traditionally gamma-ray bursts (GRBs) are classified in the T_{90} -hardness ratio two-dimensional plane into two classes: long/soft vs. short/hard GRBs. Afterglow and host galaxy observations suggest that these two categories may correspond to two different types of progenitors: those related to massive stars (Type II) and those related to compact stars (Type I). However, recent observations suggest that the duration classification scheme does not always match the physical classification scheme. Multi-wavelength criteria are needed to diagnose the physical category of a particular GRB, but some important information (e.g. supernova signature, host galaxy information) are not available during the prompt phase. In this paper, we suggest to add the “amplitude” of GRB prompt emission as the third dimension into the consideration of GRB classification with the prompt γ -ray data. We define three new parameters with the prompt emission data. The first parameter is defined as $f = F_p/F_B$, where F_p and F_B are the peak flux and the background flux, respectively. The second parameter is defined by $f_{\text{eff}} = F'_p/F_B$, where F'_p the peak flux of the a pseudo-GRB, which is generated by scaling the original lightcurve down until the measured T_{90} above the background is just less than 2 s. This is an “effective amplitude” of a real long GRB that is observed as a “short” GRB with its “tip-of-iceberg” above the background. The third parameter is defined as $f_{\text{eff},z} = F'_{p,z}/F_B$, where $F'_{p,z}$ is the peak flux of another pseudo-GRB, which is generated by progressively moving a GRB to a higher redshift z , until its measured “rest frame” duration $T_{90}/(1+z)$ is just below 2 s. We systematically derive these parameters for the *Swift* GRBs detected between December 2004 to December 2011. We find that most short GRBs are likely not “tip-of-iceberg” of long GRBs. However, one needs to be cautious if a short GRB has $f < 2$, since it could be either intrinsically short or intrinsically long with the extended emission episodes buried beneath the background. We also find that most long GRBs would show up as rest-frame short GRBs if they have a high enough redshift. This interprets the interesting observational fact that the three highest- z GRBs (080913, 090423, 090429B) all have a rest frame duration shorter than 2 s, but are all likely Type II GRBs.

Subject headings: gamma-ray bursts: general — methods: statistical

1. INTRODUCTION

Traditionally, gamma-ray bursts (GRBs) are classified based on duration (T_{90}) and hardness ratio (HR) of their prompt gamma-ray emission. In the CGRO/BATSE era, GRBs were classified into two categories in the T_{90} -HR two-dimensional plane (Kouveliotou et al. 1993) with a rough separation in the duration dimension at $T_{90} \sim 2$ s. Long GRBs are typically soft while short GRBs are typically hard, so that the two classes cluster in two regions in the T_{90} -HR plane. Such a distribution is energy-dependent and instrument-dependent (e.g. Qin et al. 2012; Zhang et al. 2012).

Broad-band afterglow observations of long GRBs reveal that their host galaxies are typically irregular galaxies with intense star formation (Fruchter et al. 2006). Some long GRBs are firmly associated with Type Ib/c supernova (e.g. Hjorth et al. 2003; Stanek et al. 2003; Campana et al. 2006; Pian et al. 2006). This strongly

suggests that they are likely related to the deaths of massive stars, and the “collapsar” model has been widely accepted to be the standard paradigm for long GRBs (Woosley 1993; MacFadyen & Woosley 1999). Detections of afterglows and host galaxies of short GRBs in the *Swift* (Gehrels et al. 2004) era have advanced our understanding of their physical origin. Some short GRBs are found to be associated with nearby early-type galaxies with little star formation (Gehrels et al. 2005; Bloom et al. 2006; Barthelmy et al. 2005; Berger et al. 2005) or have a large offset from the host even if they are associated with star forming galaxies (e.g. Fox et al. 2005; Fong et al. 2010). Deep upper limits of their supernova signals are obtained (Kann et al. 2011 and references therein). This points towards an origin that does not involve a massive star. The leading scenario is mergers of two neutron stars (Paczýnski 1986; Eichler et al. 1989) or mergers of a neutron star and a black hole (Paczýnski 1991).

Further observations revealed a more complicated picture, suggesting that duration is no longer a reliable indicator of the physical origin of a GRB. The detections of two nearby long-duration GRBs without association of a supernova, i.e. GRB 060614 ($T_{90} \sim 100$ s at $z = 0.125$) and GRB 060505 ($T_{90} = 4$ s at $z = 0.089$, cast doubts on that all long GRBs are of a massive star origin (Gehrels et al. 2006; Gal-Yam et al. 2006; Fynbo et al. 2006;

¹ Department of Physics and Astronomy, University of Nevada, Las Vegas, NV 89154, USA; lhj@physics.unlv.edu, zhang@physics.unlv.edu.

² Department of Physics and GXU-NAOC Center for Astrophysics and Space Sciences, Guangxi University, Nanning 530004, China

³ Department of Astronomy and Astrophysics, The Pennsylvania State University, University Park, PA 16802, USA

⁴ Department of Physics and Mathematics, Aoyama Gakuin University, Sagami-hara-shi Kanagawa 252-5258, Japan

Della Valle et al. 2006). On the other hand, some properties of GRB 060614 (e.g. short spectral lag, Gehrels et al. 2006) and the large offset from the star forming region in the host (Gal-Yam et al. 2006) are consistent with being a compact star origin. Zhang et al. (2007b) showed that if GRB 060614 were somewhat less energetic, it would appear as quite similar to GRB 050724, which is the “smoking gun” short GRB (with extended emission) that suggests a compact star origin (Barthelmy et al. 2005; Berger et al. 2005). Later, several high- z GRBs with the rest frame duration $T_{90}/(1+z)$ shorter than 2 s were discovered: GRB 080913 at $z = 6.7$ with $T_{90} = 8$ s (Greiner et al. 2009), GRB 090423 at $z = 8.2$ with $T_{90} = 10.3$ s (Tanvir et al. 2009; Salvaterra et al. 2009), and GRB 090429B at $z = 9.4$ with $T_{90} = 5.5$ s (Cucchiara et al. 2011), but various arguments suggest that they are of a massive star origin (Zhang et al. 2009). Later, more traditional short GRBs are found to be likely of a massive star origin. For example, GRB 090426, at $z = 2.609$, is found to have an observed BAT band duration $T_{90} = 1.2 \pm 0.3$ s and a rest frame duration $T_{90}/(1+z) \sim 0.33$ s, but its other properties are fully consistent with being of a massive star origin (Levesque et al. 2010; Xin et al. 2011; Thöne et al. 2011).

In view of these complications, Zhang (2006) and Zhang et al. (2007b) suggested to classify GRBs physically into Type II (massive star origin) and Type I (compact star origin). Zhang et al. (2009) studied the statistical properties of the Type II and Type I Gold Samples, and found that although the Type II Gold sample tracks the bulk of long GRBs well, the Type I Gold sample is not a good representative of the short GRBs. They suggested a set of multi-wavelength criteria to diagnose the physical origin of GRBs (see also Kann et al. 2011), and suspected that some, maybe most high-redshift high-luminosity short GRBs are probably of a Type II origin. This conclusion was later also drawn by several groups independently based on very different arguments (Virgili et al. 2011; Cui et al. 2012; Bromberg et al. 2012).

Even though the multi-wavelength criteria can give more definite clues about the origin of a GRB, they are not available promptly after the trigger of the GRB. Some criteria that carry most weight (e.g. supernova signature, host galaxy information) need late, deep optical observations. It is still useful to apply the prompt gamma-ray data to dig out more information, which may be helpful to infer the physical origin of a GRB. For example, in Lü et al. (2010), we have proposed a new observational parameter ε defined by $E_{\gamma,iso}/E_{p,z}^{5/3}$, where $E_{\gamma,iso}$ is the burst isotropic gamma-ray energy and $E_{p,z}$ is the rest-frame spectral peak energy. This parameter has a cleaner bimodal distribution, and the two types of burst classified with the ε criterion match the physical classification scheme (Type I vs. Type II) better. This method still needs the redshift information.

In this paper, we propose to add a third dimension “amplitude” into consideration to classify GRBs using the prompt gamma-ray data (see a preliminary discussion in Zhang 2012). The motivation is to study the possibility that a real long GRB may be observed as a “short” one if the majority of emission episode is too faint to be detected above the background. We call this the “tip-of-iceberg” effect. To quantify this effect, in Sect. 2,

we introduce a new “amplitude parameter” f , and study the distribution of *Swift* GRBs in the three-dimensional ($T_{90} - \text{HR} - f$) space. In Sect. 3, we introduce an effective amplitude parameter f_{eff} to discuss the range of amplitude if a long GRB is observed as “short” due to the tip-of-iceberg effect. We compare the range of f distribution of short GRBs and the f_{eff} distribution of long GRBs and suggest a confusion regime of f where an observed short GRB may be in fact long. In Sect. 4, we define a parameter $f_{\text{eff},z}$ by “moving” GRBs with known redshift to higher redshifts through simulations until they become “rest-frame short” GRBs. We take GRB 080319B as an example, and show that long GRBs can become rest-frame short GRBs at high enough redshifts, but with a moderately large f . We show that this is consistent with the three highest- z GRBs: 080913, 090423 and 090429B. We draw conclusions in Sect. 5 with some discussion.

2. THE AMPLITUDE PARAMETER

In the previous T_{90} -HR two-dimensional diagram, the amplitude information of GRBs is missing. Some GRBs can be very bright, while some others can be faint and barely above the threshold. A bright burst can have more emission episodes emerge above the background, so for a same observed T_{90} , a fainter burst may be intrinsically longer than a brighter burst. So this third dimension, i.e. the “amplitude”, carries important information and should be introduced in GRB classification studies.

We define the amplitude parameter as

$$f \equiv \frac{F_p}{F_B}, \quad (1)$$

where F_p is the 1-second peak flux on the gamma-ray emission lightcurve, and F_B is the average background flux of the burst. Both fluxes are in units of count rate.

We systematically process the *Swift* Burst Alert Telescope (BAT) GRB data to extract lightcurves. We developed an IDL script to automatically download and maintain all the *Swift* BAT data. We use the standard HEASoft tools (version 6.12) to process the data. By running *bateconvert* from the HEASoft software release, we obtain the energy scale for the BAT events. The lightcurves are extracted by running *batbinevt* (Sakamoto et al. 2007). For each burst, we calculate the cumulative distribution of the source count using the arrival times of a fraction of 5% and 95% of the total counts to define T_{90} . The time bin size is fixed to 64 ms for all the bursts.

Our sample includes the GRBs detected by *Swift* BAT from December 2004 to December 2011. We only selected 437 GRBs with S/N ratio higher than 5, which include 395 long GRBs and 42 short GRBs. Among them, 182 have redshift measurements. For each GRB, we fit the background flux level F_B using the time intervals before and after the burst. This background is burst-dependent, but is around a value of 8000 cts/s. For a small fraction (6.8%) of the bursts, the background before and after the burst is uneven. This is because some bright hard X-ray sources could be entering or exiting the BAT field of view during the slew. For these cases, we fit the background before and after the burst with a straight line with a slope. F_B is defined by fitted background flux at the peak time⁵. Figure 1 shows the histogram of F_B for all

⁵ This flux level is usually slightly higher than the “true” back-

the GRBs in our sample.

The f values of the GRBs in our sample are presented in our group website <http://grb.physics.unlv.edu/f/data.txt>. The $T_{90} - \text{HR} - f$ 3-dimensional distribution diagram of *Swift* GRBs is shown in Fig.2. Long and short GRBs are denoted as black and white symbols. The projections in the $T_{90} - \text{HR}$, $T_{90} - f$ and $\text{HR} - f$ planes are denoted in red, green, and blue colors, respectively, with long and short GRBs denoted by the filled and open symbols, respectively. In Fig.3a, we show the $T_{90} - f$ diagram with different symbols denoting different types of GRBs: gray for long GRBs, red for short GRBs, blue for short GRBs with extended emission (T_{90} calculated by excluding the extended emission), magenta for the three “rest-frame short” ($T_{90}/(1+z) < 2$ s) high- z GRBs, black for other “rest-frame short” GRBs, and two special GRBs, 090426 and 060614, are marked separately.

The distributions of the f -parameter for both long and short GRBs are presented in Fig.4. As expected, Most bursts are clustered around small f values, and only a small fraction of bursts have $f > 3$. The f distribution can be roughly fit as a power law function, i.e. $N(f) \propto f^{-a}$, with $a \sim 3.15$ for long GRBs and $a \sim 1.5$ for short GRBs. The mean value of f is $\bar{f} = 1.48$ for long GRBs and $\bar{f} = 1.82$ for short GRBs. The largest f values for both long and short GRBs are around 10. The relative paucity of small f for short GRBs may be understood as a selection effect (Sakamoto et al. 2008, 2011): Short GRBs are detected via “rate triggers”, which require a relatively large f value to meet the trigger criterion. On the other hand, long GRBs can be caught via “imaging triggers” near the threshold, so that they can be detected with lower f values close to unity.

3. EFFECTIVE AMPLITUDE OF LONG GRBS TO BE CONFUSED AS SHORT GRBS

A long GRB may be confused as a short GRB if only its brightest spikes with duration shorter than 2 s are above the background. To quantify such a tip-of-iceberg effect, we define an “effective amplitude” of a GRB as

$$f_{\text{eff}} \equiv \frac{F'_p}{F_B}. \quad (2)$$

Here F'_p is the 1-second peak flux of a pseudo GRB, which is linearly scaled down from an original GRB until its signal above the background has a duration $T_{90,\text{eff}}$ just shorter than 2 s. The physical meaning of the pseudo GRB is an otherwise identical GRB at the same redshift, except that the amplitude is lower by a factor f/f_{eff} . For a short GRB, f_{eff} is identical to f since the signal is already shorter than 2 s in the original burst. Figure 5 shows the case of the long GRB 050525A as an example. The original burst (black curve) has an $f = 9.43$. After linearly scaling it down by a factor of 2.44 on the y-axis, most of the signal is buried below the background level F_B (the gray curve, with the sum of signal and background being at most the background level), so that the signal above the background (red curve) is just shorter

ground level due to the source contamination. However, this is not a concern for our analysis, since we are investigating the tip-of-iceberg effect with respect to the background at the detection time.

than 2 s. The amplitude of this pseudo burst is the effective amplitude of the original burst, which is $f_{\text{eff}} = 3.86$ for GRB 050525A.

Figure 3b gives the $T_{90} - f_{\text{eff}}$ distribution of all the GRBs in our sample. The short GRBs have the same distribution as f , while long GRBs have systematically smaller values of f_{eff} than f . The f_{eff} distribution histogram of long GRBs is also shown in Fig.4 (blue dots line), which has a mean value $\bar{f}_{\text{eff}} = 1.28$, and can be fit with a power law with $a = 4.23 \pm 0.92$.

One immediate conclusion from Fig.3b and Fig.4 is that the distribution of f_{eff} of long GRBs is very different from the f distribution of short GRBs. This suggests that the majority of short GRBs are *not* tip-of-iceberg of long GRBs. Instead, they reflect the intrinsically short duration of the central engine.

Nonetheless, Fig.3b raises a caution for those detected short GRBs with $f < 2$. Even though most of them may be still intrinsically short, it is possible that some of them may be tip-of-iceberg of long GRBs. This is because a non-negligible fraction of long GRBs have f_{eff} falling into this range. The high- z short GRB 090426 (Levesque et al. 2010; Xin et al. 2011; Thöne et al. 2011) has $f = 1.48$, which falls right in this range.

On the other hand, GRB 060614 (Gehrels et al. 2006), the long GRB without supernova association, has $f_{\text{eff}} = 2.26$. Its tip-of-iceberg still has a large f to be consistent with the short GRB f distribution. Indeed, by scaling it down, it looks like a short GRB with extended emission (Zhang et al. 2007b). Our analysis again supports the Type I (compact star) origin of this GRB.

4. “REST-FRAME SHORT” GRBS

Some long GRBs have a rest-frame duration $T_{90}/(1+z) < 2$ s. The three GRBs with the highest redshifts, i.e. GRB 080913 (Greiner et al. 2009), GRB 090423 (Tanvir et al. 2009; Salvaterra et al. 2009), and GRB 090429B (Cucchiara et al. 2011) are all of this type, but likely have a Type II (massive star) origin based on the multi-wavelength criteria (Zhang et al. 2009). It would be very interesting to investigate whether this is also due to the tip-of-iceberg effect.

In order to check such a possibility, we define a third parameter

$$f_{\text{eff},z} \equiv \frac{F'_{p,z}}{F_B}. \quad (3)$$

Here $F'_{p,z}$ is the 1-second peak flux of a pseudo GRB, which is generated by moving the original GRB to progressively higher redshifts until the rest-frame duration $T_{90}/(1+z)$ becomes shorter than 2 s. A GRB, when moved to a higher redshift, would usually have a shorter rest frame duration, although the observed duration may not shrink due to time dilation (Kocevski & Petrosian 2011). In principle, it would always reach the “rest-frame-short” phase before completely disappearing beneath the background. It would be interesting to investigate the critical redshift z_c above which a burst appears as rest-frame-short.

Technically, moving a GRB with known redshift to higher redshifts is not straightforward. One needs to reduce the time-resolved spectra of the GRB, derive the correct spectral parameters, and perform a proper k -

correction to the spectrum in order to obtain the BAT-band light curve of the pseudo GRB.

To carry out such an exercise, for each GRB with redshift measurement, we first apply *Xspec* to conduct a time-dependent spectral analysis to the raw data. We dissect the lightcurve into multiple time bins, with the bin size self-adjusted to allow a signal-to-noise ratio $S/N > 5$, so that the spectral parameters can be constrained. A typical GRB spectrum, if the observational band is wide enough, can be described as the Band function (Band et al. 1993; Abdo et al. 2009; Zhang et al. 2011). In order to perform a proper k correction, ideally one should know the Band spectral parameters α , β and E_p . However, since the BAT band is narrow, for most GRBs the spectra can be only fit by a cutoff power law or a single power law (Sakamoto et al. 2008, 2011). We therefore apply the following procedure to estimate the Band spectral parameters: 1. If a burst was also detected by *Fermi* GBM or *Konus* Wind, we adopt the spectral parameters measured by those instruments. 2. For those bursts that are not detected by other instruments but can be fit with a cutoff power law, we adopted the derived α and E_p parameters, and assume a typical value $\beta = -2.3$. 3. For those GRBs that can only be fit with a single power law, we have to derive E_p using an empirical correlation between the BAT-band photon index Γ^{BAT} and E_p , as derived previously for Swift GRBs (Sakamoto et al. 2009; Zhang et al. 2007a,b; Virgili et al. 2011). The typical parameters $\alpha = -1$, $\beta = -2.3$ are adopted to perform the simulations. We note that moving a GRB to a higher z is effectively observing the rest-frame spectra in a higher energy band given the same observed BAT band. The spectral parameters β and E_p are therefore essential. These parameters are unfortunately usually not available for the *Swift* GRBs. So our pseudo GRBs should be considered only as simulated GRBs rather than the original GRBs being moved to higher redshifts. In any case, such a simulation can serve the purpose of investigating the tip-of-iceberg selection effect. A similar simulation was carried out by Kocevski & Petrosian (2011).

Given the spectral parameters α , β and E_p of a particular GRB with known redshift z , we use the following procedure to simulate the pseudo GRB. First, we calculate the time-dependent bolometric burst luminosity using

$$L(t) = 4\pi D_L^2(z) F(t) k, \quad (4)$$

where $F(t)$ is the BAT-band, time-dependent flux, $D_L(z)$ is the luminosity distance to the source at the redshift z , and the k -correction factor corrects the BAT-band (15 – 150 keV) flux to a wide band in the burst rest frame ($1 - 10^4$ keV in this analysis), i.e.

$$k = \frac{\int_{1/(1+z)}^{10^4/(1+z)} E N(E) dE}{\int_{15}^{150} E N(E) dE}. \quad (5)$$

Here $N(E)$ is the time-dependent Band photon spectrum. To calculate $D_L(z)$, the concordance cosmology parameters ($H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.30$, and $\Omega_\Lambda = 0.70$) are adopted.

Next, we apply the spectral model to calculate the BAT-band flux for a pseudo GRB at redshift z' . We

keep the bolometric luminosity as a constant, and derive the BAT band flux using

$$F'(t') = \frac{L(t)}{4\pi D_L^2(z') k'}, \quad (6)$$

where

$$t' = \frac{1+z'}{1+z} t, \quad (7)$$

$D_L(z')$ is the luminosity distance to the source at redshift z' , and

$$k' = \frac{\int_{1/(1+z')}^{10^4/(1+z')} E' N(E') dE'}{\int_{15}^{150} E' N(E') dE'}. \quad (8)$$

Here $N(E')$ is the observed photon number spectrum of the pseudo GRB. The spectrum is still a Band function with the same α and β values. The only difference is that the peak energy is now shifted to $E'_p = E_p(1+z)/(1+z')$. We then add the background level F_B to the signal, and re-calculate T_{90} of the pseudo GRB for each z' , and calculate the rest-frame duration $T_{90}/(1+z)$. By progressively increasing z' , we identify a critical redshift z_c beyond which $T_{90}/(1+z) < 2$ s is satisfied. The peak flux of the pseudo GRB at z_c is used to define $f_{\text{eff},z}$. We then continue to increase the redshift, until the entire GRB disappears below the background. We record this redshift as z_{max} . The redshift range (z_c, z_{max}) is then where a rest-frame short GRB can be observed.

As an example, we take the “naked-eye” GRB 080319B (Racusin et al. 2008) as the original burst and perform the simulation. The results are shown in Fig.6. The time-integrated γ -ray spectrum is well fit using a Band function, with $E_p = 675 \pm 22$ keV. The time-resolved of spectra are well obtained, with E_p evolving from ~ 740 keV to ~ 540 keV. The rest-frame isotropic energy release is $E_{\text{iso}} = (1.14 \pm 0.09) \times 10^{54}$ erg in the source frame $1 - 10^4$ keV band (Racusin et al. 2008, Amati et al. 2008).

We apply the above method to simulate pseudo GRBs with increasing redshifts. The lightcurves of the pseudo GRBs are presented in Fig.6a. Different colors denote different redshifts. From top to bottom, the redshifts are: $z = 0.937$ (original), 1, 2.3, 2.8, 3.6, 4.5, 5.1, and the critical redshift is $z_c = 5.53$. As shown in Fig.6a, the peak flux of the pseudo GRBs become progressively lower as z increases, and the observed durations initially become longer (due to time dilation) but later shrink (due to tip-of-iceberg effect). The rest-frame duration $T_{90}/(1+z)$ is found to decrease with redshift, similar to track with a smooth broken power-law (Fig.6b). At $z = z_c = 5.53$, $T_{90}/(1+z)$ becomes shorter than 2 s. We derive $f_{\text{eff},z} = 1.41$. The burst is no longer detectable at $z = z_{\text{max}} = 5.92$.

We carry out the same exercise for all the *Swift* GRBs with known redshifts. The $T_{90}/(1+z) - f_{\text{eff},z}$ diagram is presented in Fig.3c. We can see that $f_{\text{eff},z}$ are all below ~ 1.7 . It is interesting to note that the three highest- z GRBs (080913, 090423 and 090429B) and other rest-frame-short GRBs all have f values within this range. This suggests that they are simply the tip-of-iceberg of long GRBs. This conclusion is consistent with their Type II origin as derived from multi-wavelength arguments (Zhang et al. 2009). In Fig.7, we plot the histograms of z_c and z_{max} of all the GRBs in our analysis, and compare

them with the z distribution of the detected rest-frame short GRBs. It is found that they are generally consistent with each other. The discrepancy in the high- z end (the distribution does not fully include the highest z GRB) may be due to the uncertainty of the high-energy spectra used in our simulations.

If the rest-frame short GRBs are the tip-of-iceberg of long GRBs, then the extended emission episodes (“icebergs” themselves) may show up in the softer X-ray band. To test this possibility, in Fig.8 we simulate the expected XRT band lightcurve of a pseudo naked-eye GRB 080319B at $z = z_c = 5.53$ (black). The same k -correction method has been applied. This is compared against the XRT-band lightcurves of the three highest- z GRBs (green for GRB 080913, blue for GRB 090423, and red for GRB 090429B), as well as the original XRT-band data of GRB 080319B (gray). It is seen that the XRT lightcurve of the pseudo GRB has an extended flaring episode extending to ~ 200 s followed by a steep decay, which is similar to the case of GRB 090423.

5. CONCLUSIONS AND DISCUSSION

In this paper, we propose to add “amplitude” as the third dimension to study GRB classification using the prompt emission data. We introduced three parameters, f (Eq.(1)), f_{eff} (Eq.(2)), and $f_{\text{eff},z}$ (Eq.(3)), to describe the amplitude of the original GRB and some simulated pseudo GRBs. We find the following interesting results:

- The f parameters for both long and short GRBs are distributed between 1 and about 10 as a rough power law. The paucity of low- f short GRBs may be understood as a trigger selection effect.
- The f parameter of many short GRBs is larger than the f_{eff} parameter of long GRBs. This suggests that most short GRBs are likely intrinsically short, and *not* simply the tip-of-iceberg of long GRBs.

- There is a confusion regime at $f < 2$ for short GRBs, since the f_{eff} parameter of a good fraction of long GRBs is in this range. It is possible that some low-amplitude ($f < 2$) short GRBs can be in fact “long”. This is indeed the case of the Type-II short GRB 090426. Most low-amplitude short GRBs may be still intrinsically short. Nonetheless, the possibility of contamination from long GRBs is not small. We suggest that in the future, one should be cautious to classify short GRBs with $f < 2$.
- When long GRBs are moved to high redshifts, they are likely observed as rest-frame short GRBs due to the “tip-of-iceberg” effect. These rest-frame short GRBs are supposed to have a low amplitude $f < 1.7$. The observed three highest- z GRBs and other rest-frame short GRBs all have such a low amplitude. So they are consistent with being tip-of-iceberg of long GRBs.

We acknowledge the use of the public data from the *Swift* data archive. This work was supported by NASA NNX10AD48G and NSF AST-0908362 (BZ). EWL acknowledges support from the National Natural Science Foundation of China under grants No. 11063001, 10873002 and 11025313, the National Basic Research Program (“973” Program) of China (Grant 2009CB824800), Special Foundation for Distinguished Expert Program of Guangxi, the Guangxi SHI-BAI-QIAN project (Grant 2007201). BBZ acknowledges support by NASA SAO G01-12102X.

REFERENCES

- Abdo, A. A., Ackermann, M., Arimoto, M., et al. 2009, *Science*, 323, 1688
- Amati, L., et al. 2008, *MNRAS*, 391, 577
- Band, D., Matteson, J., Ford, L., et al. 1993, *ApJ*, 413, 281
- Barthelmy, S. D., et al. 2005, *Nature*, 438, 994
- Berger, E., Kulkarni, S. R., Fox, D. B., et al. 2005, *ApJ*, 634, 501
- Bloom, J. S., Prochaska, J. X., Pooley, D., et al. 2006, *ApJ*, 638, 354
- Bromberg, O., Nakar, E., Piran, T., & Sari, R. 2012, *arXiv:1210.0068*
- Campana, S., Mangano, V., Blustin, A. J., et al. 2006, *Nature*, 442, 1008
- Cucchiara, A., Levan, A. J., Fox, D. B., et al. 2011, *ApJ*, 736, 7
- Cui, X.-H., Nagataki, S., Aoi, J., & Xu, R.-X. 2012, *Research in Astronomy and Astrophysics*, 12, 1255
- Della Valle, M., Chincarini, G., Panagia, N., et al. 2006, *Nature*, 444, 1050
- Eichler, D., Livio, M., Piran, T., & Schramm, D. N. 1989, *Nature*, 340, 126
- Fong, W., Berger, E., & Fox, D. B. 2010, *ApJ*, 708, 9
- Fox, D. B., Frail, D. A., Price, P. A., et al. 2005, *Nature*, 437, 845
- Fruchter, A. S., Levan, A. J., Strolger, L., et al. 2006, *Nature*, 441, 463
- Fynbo, J. P. U., Watson, D., Thöne, C. C., et al. 2006, *Nature*, 444, 1047
- Gal-Yam, A., Fox, D. B., Price, P. A., et al. 2006, *Nature*, 444, 1053
- Gehrels, N., Chincarini, G., Giommi, P., et al. 2004, *ApJ*, 611, 1005
- Gehrels, N., Norris, J. P., Barthelmy, S. D., et al. 2006, *Nature*, 444, 1044
- Gehrels, N., Sarazin, C. L., O’Brien, P. T., et al. 2005, *Nature*, 437, 851
- Greiner, J., Krühler, T., Fynbo, J. P. U., et al. 2009, *ApJ*, 693, 1610
- Hjorth, J., Sollerman, J., Møller, P., et al. 2003, *Nature*, 423, 847
- Kann, D. A., Klose, S., Zhang, B., et al. 2011, *ApJ*, 734, 96
- Kocevski, D., & Petrosian, V. 2011, *arXiv:1110.6175*
- Kouveliotou, C., Meegan, C. A., Fishman, G. J., et al. 1993, *ApJ*, 413, L101
- Lü, H.-J., Liang, E.-W., Zhang, B.-B., & Zhang, B. 2010, *ApJ*, 725, 1965
- Levesque, E. M., Bloom, J. S., Butler, N. R., et al. 2010, *MNRAS*, 401, 963
- MacFadyen, A. I., & Woosley, S. E. 1999, *ApJ*, 524, 262
- Paczynski, B. 1986, *ApJ*, 308, L43
- Paczynski, B. 1991, *Acta Astron* 41, 257
- Pian, E., Mazzali, P. A., Masetti, N., et al. 2006, *Nature*, 442, 1011
- Qin, Y., Liang, E.-W., Yi, S.-X., et al. 2012, *arXiv:1205.1188*
- Racusin, J. L., Karpov, S. V., Sokolowski, M., et al. 2008, *Nature*, 455, 183
- Sakamoto, T., Barthelmy, S. D., Barbier, L., et al. 2008, *ApJS*, 175, 179
- Sakamoto, T., Barthelmy, S. D., Baumgartner, W. H., et al. 2011, *ApJS*, 195, 2
- Sakamoto, T., Hill, J. E., Yamazaki, R., et al. 2007, *ApJ*, 669, 1115




FIG. 1.— The the histogram of F_B for all the GRBs in our sample.

- Sakamoto, T., Sato, G., Barbier, L., et al. 2009, *ApJ*, 693, 922
 Salvaterra, R., Della Valle, M., Campana, S., et al. 2009, *Nature*, 461, 1258
 Stanek, K. Z., Matheson, T., Garnavich, P. M., et al. 2003, *ApJ*, 591, L17
 Tanvir, N. R., Fox, D. B., Levan, A. J., et al. 2009, *Nature*, 461, 1254
 Thöne, C. C., Campana, S., Lazzati, D., et al. 2011, *MNRAS*, 414, 479
 Virgili, F. J., Zhang, B., O'Brien, P., & Troja, E. 2011, *ApJ*, 727, 109
 Woosley, S. E. 1993, *ApJ*, 405, 273
 Xin, L.-P., Liang, E.-W., Wei, J.-Y., et al. 2011, *MNRAS*, 410, 27
 Zhang, B.-B., Zhang, B., Liang, E.-W., et al. 2011, *ApJ*, 730, 141
 Zhang, B. 2012, *IAU Symposium*, 279, 102
 Zhang, B. 2006, *Nature*, 444, 1010
 Zhang, B., Liang, E., Page, K. L., et al. 2007a, *ApJ*, 655, 989
 Zhang, B., Zhang, B.-B., Liang, E.-W., et al. 2007b, *ApJ*, 655, L25
 Zhang, B., Zhang, B.-B., Virgili, F. J., et al. 2009, *ApJ*, 703, 1696
 Zhang, F.-W., Shao, L., Yan, J.-Z., & Wei, D.-M. 2012, *ApJ*, 750, 88




FIG. 2.— The $T_{90} - HR - f$ 3-dimensional distribution of the *Swift* GRBs in our sample. Long and short GRBs are denoted as solid black and white diamonds, respectively. Their projections to the $T_{90} - HR$, $T_{90} - f$, and $HR - f$ 2-dimensional planes are denoted in red, green and blue colors, respectively, with the long and short GRBs denoted with solid and open symbols, respectively.




FIG. 3.— (a): The $T_{90}-f$ diagram of the GRBs in our sample. (b) The $T_{90}-f_{\text{eff}}$ diagram of GRBs in our sample. (c) The $T_{90}/(1+z)-f_{\text{eff},z}$ diagram of GRBs in our sample. The following convention is adopted for all three plots: Gray: long GRBs, red: short GRBs; blue: short GRBs with extended emission; magenta: three GRBs with the highest z ; black: “rest-frame short” GRBs. GRB 060614 and GRB 090426 are marked with special symbols. The vertical dashed line is the 2 s separation line.




FIG. 4.— The distributions of f (for both long and short GRBs) and f_{eff} (for long GRBs only) as well as their power law fits.




FIG. 5.— An example of defining f_{eff} with GRB 050525A. The original lightcurve is shown as the black curve. The scaled-down lightcurve is shown as red above the background and gray below the background. The measured duration is shorter than 2 s (red line). The peak-to-background flux ratio of the pseudo GRB (red) is defined as f_{eff} of GRB 050525A.




FIG. 6.— (a) The simulated BAT-band pseudo GRB lightcurves by moving GRB 080319B to progressively high redshifts. From top to bottom, $z = 0.937, 1, 2.3, 2.8, 3.6, 4.5, 5.1, 5.53$. (b) The measured rest-frame duration $T_{90}/(1+z)$ of the pseudo GRBs in our simulation. The red solid line shows a smooth broken power-law fit.




FIG. 7.— The distributions of z_c and $z_{\max}$ for the simulated pseudo GRBs as compared with the redshift distribution of the observed rest-frame short GRBs.




FIG. 8.— The simulated XRT-band lightcurve of the pseudo GRB by moving GRB 080319B to $z = z_c = 5.53$ (black), as compared with the original XRT-band lightcurves of GRB 080319B (gray), GRB 080913 (green), GRB 090423 (blue), and GRB 090429B (red).